

## Medicare Chronic Condition Counts

**Table B.2.a.** Medicare beneficiary prevalence\* for 30 CCW chronic conditions\*\* using fee-for-service (FFS) claims, 2017–2021

| Totals                    | 2017 denominator <sup>#</sup> | 2017 percent | 2018 denominator <sup>#</sup> | 2018 percent | 2019 denominator <sup>#</sup> | 2019 percent |
|---------------------------|-------------------------------|--------------|-------------------------------|--------------|-------------------------------|--------------|
| Total <sup>†</sup>        | 32,483,529                    | 100%         | 32,184,841                    | 100%         | 31,860,990                    | 100%         |
| Females only <sup>‡</sup> | 17,820,358                    | 54.9%        | 17,626,850                    | 54.8%        | 17,412,231                    | 54.7%        |
| Males only <sup>▲</sup>   | 14,663,165                    | 45.1%        | 14,557,985                    | 45.2%        | 14,448,754                    | 45.3%        |

| Chronic condition**                                     | 2017 numerator <sup>#</sup> | 2017 rate | 2018 numerator <sup>#</sup> | 2018 rate | 2019 numerator <sup>#</sup> | 2019 rate |
|---|-----------------------------|-----------|-----------------------------|-----------|-----------------------------|-----------|
| Acute myocardial infarction                             | 378,696                     | 1.2%      | 391,967                     | 1.2%      | 400,692                     | 1.3%      |
| Alzheimer's disease                                     | 1,021,002                   | 3.1%      | 975,599                     | 3.0%      | 930,049                     | 2.9%      |
| Anemia  | 7,040,563                   | 21.7%     | 6,984,615                   | 21.7%     | 6,941,320                   | 21.8%     |
| Asthma  | 2,552,924                   | 7.9%      | 2,487,089                   | 7.7%      | 2,431,373                   | 7.6%      |
| Atrial fibrillation and flutter                         | 4,261,499                   | 13.1%     | 4,294,182                   | 13.3%     | 4,338,834                   | 13.6%     |
| Benign prostatic hyperplasia <sup>▲</sup>               | 3,731,131                   | 25.4%     | 3,807,277                   | 26.2%     | 3,887,623                   | 26.9%     |
| Cancer, breast (female) <sup>‡</sup>                    | 1,325,734                   | 7.4%      | 1,338,988                   | 7.6%      | 1,356,944                   | 7.8%      |
| Cancer, breast (male) <sup>▲</sup>                      | 19,570                      | 0.1%      | 25,169                      | 0.2%      | 28,217                      | 0.2%      |
| Cancer, colorectal                                      | 534,198                     | 1.6%      | 526,699                     | 1.6%      | 520,113                     | 1.6%      |
| Cancer, endometrial <sup>‡</sup>                        | 171,441                     | 1.0%      | 176,062                     | 1.0%      | 180,248                     | 1.0%      |
| Cancer, lung  | 411,784                     | 1.3%      | 414,284                     | 1.3%      | 419,811                     | 1.3%      |
| Cancer, prostate <sup>▲</sup>                           | 1,310,686                   | 8.9%      | 1,319,521                   | 9.1%      | 1,339,503                   | 9.3%      |
| Cancer, urologic (kidney, renal pelvis, and ureter)     | 241,732                     | 0.7%      | 247,779                     | 0.8%      | 255,399                     | 0.8%      |
| Cataract  | 8,388,282                   | 25.8%     | 8,371,047                   | 26.0%     | 8,364,322                   | 26.3%     |
| Chronic kidney disease                                  | 5,296,046                   | 16.3%     | 5,429,683                   | 16.9%     | 5,600,301                   | 17.6%     |
| Chronic obstructive pulmonary disease                   | 5,155,458                   | 15.9%     | 5,033,696                   | 15.6%     | 4,882,331                   | 15.3%     |
| Depression, bipolar, or other depressive mood disorders | 6,299,572                   | 19.4%     | 6,371,448                   | 19.8%     | 6,431,278                   | 20.2%     |
| Diabetes  | 9,081,954                   | 28.0%     | 8,919,842                   | 27.7%     | 8,771,519                   | 27.5%     |
| Glaucoma  | 4,483,052                   | 13.8%     | 4,545,279                   | 14.1%     | 4,615,489                   | 14.5%     |
| Heart failure and non-ischemic heart disease            | 3,940,948                   | 12.1%     | 3,965,162                   | 12.3%     | 3,987,434                   | 12.5%     |
| Hip/Pelvic fracture                                     | 377,066                     | 1.2%      | 374,278                     | 1.2%      | 374,464                     | 1.2%      |
| Hyperlipidemia  | 19,927,458                  | 61.3%     | 19,967,870                  | 62.0%     | 20,040,500                  | 62.9%     |
| Hypertension  | 22,024,151                  | 67.8%     | 21,801,519                  | 67.7%     | 21,592,541                  | 67.8%     |
| Hypothyroidism  | 6,788,655                   | 20.9%     | 6,803,999                   | 21.1%     | 6,802,822                   | 21.4%     |
| Ischemic heart disease                                  | 7,081,526                   | 21.8%     | 7,000,441                   | 21.8%     | 6,949,697                   | 21.8%     |

| Chronic condition**                                | 2017 numerator <sup>#</sup> | 2017 rate | 2018 numerator <sup>#</sup> | 2018 rate | 2019 numerator <sup>#</sup> | 2019 rate |
|--|-----------------------------|-----------|-----------------------------|-----------|-----------------------------|-----------|
| Non-Alzheimer's dementia                           | 2,587,487                   | 8.0%      | 2,529,061                   | 7.9%      | 2,474,287                   | 7.8%      |
| Osteoporosis with or without pathological fracture | 3,209,888                   | 9.9%      | 3,265,644                   | 10.1%     | 3,351,209                   | 10.5%     |
| Parkinson's disease and secondary Parkinsonism     | 553,944                     | 1.7%      | 553,460                     | 1.7%      | 556,085                     | 1.7%      |
| Pneumonia, all-cause                               | 1,779,648                   | 5.5%      | 1,736,215                   | 5.4%      | 1,682,695                   | 5.3%      |
| Rheumatoid arthritis/Osteoarthritis                | 11,524,987                  | 35.5%     | 11,552,428                  | 35.9%     | 11,588,956                  | 36.4%     |
| Stroke / Transient Ischemic Attack                 | 2,133,129                   | 6.6%      | 2,099,102                   | 6.5%      | 2,075,437                   | 6.5%      |

\* Includes all Medicare beneficiaries who were eligible for or enrolled in Medicare on or after January 1, YYYY. The Chronic Conditions Warehouse (CCW) team calculates period prevalence for these rates: beneficiaries with full or nearly full FFS coverage during the year who had claim(s) for the condition within the condition-specified look back period (30 CCW Chronic Conditions have a one- to two-year look-back period). Refer to the algorithm specifications on the [Chronic Conditions page](#) of the CCW website.

\*\* The CCW team may have counted beneficiaries in more than one chronic condition category.

† Includes beneficiaries with full or nearly full FFS coverage (i.e., 11 or 12 months of Medicare Parts A and B [or coverage until time of death] and one month or less of health maintenance organization [HMO] coverage).

‡ Includes only females for cancer, endometrial and female breast.

▲ Includes only males for benign prostatic hyperplasia and cancer, prostate and cancer, male breast.

# The CCW team defines the numerator as evidence of claims for the condition.

**NOTE:** The application of different Medicare coverage restrictions/criteria will impact the size of these samples.

**Table B.2.a.** Medicare beneficiary prevalence\* for 30 CCW chronic conditions\*\* using fee-for-service (FFS) claims, 2017–2021

| Totals                    | 2020 denominator <sup>#</sup> | 2020 percent | 2021 denominator <sup>#</sup> | 2021 percent |
|---------------------------|-------------------------------|--------------|-------------------------------|--------------|
| <b>Total†</b>             | <b>31,187,299</b>             | <b>100%</b>  | <b>29,757,982</b>             | <b>100%</b>  |
| Females only <sup>‡</sup> | 17,027,561                    | 54.6%        | 16,255,376                    | 54.6%        |
| Males only <sup>▲</sup>   | 14,159,734                    | 45.4%        | 13,502,599                    | 45.4%        |

| Chronic condition**                                     | 2020 numerator <sup>#</sup> | 2020 rate | 2021 numerator <sup>#</sup> | 2021 rate |
|---|-----------------------------|-----------|-----------------------------|-----------|
| Acute myocardial infarction                             | 384,185                     | 1.2%      | 381,206                     | 1.3%      |
| Alzheimer's disease                                     | 857,109                     | 2.7%      | 750,273                     | 2.5%      |
| Anemia  | 6,645,691                   | 21.3%     | 6,266,946                   | 21.1%     |
| Asthma  | 2,314,636                   | 7.4%      | 2,150,373                   | 7.2%      |
| Atrial fibrillation and flutter                         | 4,249,615                   | 13.6%     | 4,088,560                   | 13.7%     |
| Benign prostatic hyperplasia <sup>▲</sup>               | 3,822,081                   | 27.0%     | 3,724,246                   | 27.6%     |
| Cancer, breast (female) <sup>‡</sup>                    | 1,325,827                   | 7.8%      | 1,286,577                   | 7.9%      |
| Cancer, breast (male) <sup>▲</sup>                      | 29,966                      | 0.2%      | 31,593                      | 0.2%      |
| Cancer, colorectal                                      | 488,814                     | 1.6%      | 455,757                     | 1.5%      |
| Cancer, endometrial <sup>‡</sup>                        | 174,721                     | 1.0%      | 168,734                     | 1.0%      |
| Cancer, lung  | 405,281                     | 1.3%      | 385,721                     | 1.3%      |
| Cancer, prostate <sup>▲</sup>                           | 1,315,058                   | 9.3%      | 1,277,012                   | 9.5%      |
| Cancer, urologic (kidney, renal pelvis, and ureter)     | 252,697                     | 0.8%      | 245,085                     | 0.8%      |
| Cataract  | 6,947,683                   | 22.3%     | 7,735,428                   | 26.0%     |
| Chronic kidney disease                                  | 5,589,772                   | 17.9%     | 5,383,435                   | 18.1%     |
| Chronic obstructive pulmonary disease                   | 4,437,223                   | 14.2%     | 3,883,186                   | 13.0%     |
| Depression, bipolar, or other depressive mood disorders | 6,206,012                   | 19.9%     | 5,790,493                   | 19.5%     |
| Diabetes  | 8,416,991                   | 27.0%     | 7,873,257                   | 26.5%     |
| Glaucoma  | 4,449,662                   | 14.3%     | 4,255,705                   | 14.3%     |
| Heart failure and non-ischemic heart disease            | 3,836,105                   | 12.3%     | 3,593,979                   | 12.1%     |
| Hip/Pelvic fracture                                     | 341,689                     | 1.1%      | 336,351                     | 1.1%      |
| Hyperlipidemia  | 19,570,280                  | 62.8%     | 18,897,422                  | 63.5%     |
| Hypertension  | 20,840,730                  | 66.8%     | 19,734,885                  | 66.3%     |
| Hypothyroidism  | 6,597,347                   | 21.2%     | 6,296,573                   | 21.2%     |
| Ischemic heart disease                                  | 6,681,009                   | 21.4%     | 6,303,076                   | 21.2%     |
| Non-Alzheimer's dementia                                | 2,374,413                   | 7.6%      | 2,179,461                   | 7.3%      |
| Osteoporosis with or without pathological fracture      | 3,270,710                   | 10.5%     | 3,224,005                   | 10.8%     |

Chronic Conditions Warehouse (CCW) — Your source for national CMS Medicare and Medicaid research data

| Chronic condition**                            | 2020 numerator <sup>#</sup> | 2020 rate | 2021 numerator <sup>#</sup> | 2021 rate |
|--|-----------------------------|-----------|-----------------------------|-----------|
| Parkinson's disease and secondary Parkinsonism | 538,220                     | 1.7%      | 509,189                     | 1.7%      |
| Pneumonia, all-cause                           | 1,710,025                   | 5.5%      | 1,600,855                   | 5.4%      |
| Rheumatoid arthritis/Osteoarthritis            | 10,948,670                  | 35.1%     | 10,308,821                  | 34.6%     |
| Stroke / Transient Ischemic Attack             | 1,869,620                   | 6.0%      | 1,853,895                   | 6.2%      |

\* Includes all Medicare beneficiaries who were eligible for or enrolled in Medicare on or after January 1, YYYY. The Chronic Conditions Warehouse (CCW) team calculates period prevalence for these rates: beneficiaries with full or nearly full FFS coverage during the year who had claim(s) for the condition within the condition-specified look back period (30 CCW Chronic Conditions have a one- to two-year look-back period). Refer to the algorithm specifications on the [Chronic Conditions page](#) of the CCW website.

\*\* The CCW team may have counted beneficiaries in more than one chronic condition category.

† Includes beneficiaries with full or nearly full FFS coverage (i.e., 11 or 12 months of Medicare Parts A and B [or coverage until time of death] and one month or less of health maintenance organization [HMO] coverage).

‡ Includes only females for cancer, endometrial and female breast.

▲ Includes only males for benign prostatic hyperplasia and cancer, prostate and cancer, male breast.

# The CCW team defines the numerator as evidence of claims for the condition.

**NOTE:** The application of different Medicare coverage restrictions/criteria will impact the size of these samples.